



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 487/20, 471/20, 471/04, A61K 31/505, 31/445		A1	(11) International Publication Number: WO 99/01455
			(43) International Publication Date: 14 January 1999 (14.01.99)
(21) International Application Number: PCT/SE98/01206 (22) International Filing Date: 22 June 1998 (22.06.98) (30) Priority Data: 9702534-0 1 July 1997 (01.07.97) SE (71) Applicant (for all designated States except US): ASTRA PHARMACEUTICALS LTD. [GB/GB]; Home Park, Kings Langley, Herts WD4 8DH (GB). (71) Applicant (for MG only): ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): HAMLEY, Peter [GB/GB]; Astra Charnwood, Bakewell Road, Loughborough, Leics LE11 5RH (GB). PIMM, Austen [GB/GB]; Astra Charnwood, Bakewell Road, Loughborough, Leics LE11 5RH (GB). TINKER, Alan [GB/GB]; Astra Charnwood, Bakewell Road, Loughborough, Leics LE11 5RH (GB). (74) Agent: ASTRA AKTIEBOLAG; Patent Dept., S-151 85 Södertälje (SE).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: COMPOUNDS			
<div style="text-align: center;"> <p>(I)</p> </div>			
(57) Abstract			
<p>There are provided novel compounds of formula (I) wherein A represents an aromatic carbocyclic ring or a 5- or 6-membered heterocyclic aromatic ring containing 1 to 3 heteroatoms which may be the same or different and are selected from O, N and S; X represents $-(CH_2)_n-$ wherein n represents zero or 1; and R¹, R² and R³ are as defined in the specification, and pharmaceutically acceptable salts thereof, and enantiomers and tautomers thereof; together with processes for their preparation, compositions containing them and their use in therapy. The compounds are inhibitors of nitric oxide synthase and are thereby particularly useful in the treatment or prophylaxis of inflammatory disease and pain.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

COMPOUNDS


Field of the Invention

The present invention relates to novel compounds which are aminospiriperidine quinazoline derivatives. The invention also concerns related aspects including processes
5 for the preparation of the compounds, compositions containing them and their use as pharmaceuticals. There are also provided chemical intermediates useful for the production of the compounds.

Background of the Invention

10 Nitric oxide is produced in mammalian cells from L-arginine by the action of specific nitric oxide synthases (NOSs). These enzymes fall into two distinct classes - constitutive NOS (cNOS) and inducible NOS (iNOS). At the present time, two constitutive NOSs and one inducible NOS have been identified. Of the constitutive NOSs, an endothelial enzyme (ecNOS) is involved with smooth muscle relaxation and the regulation of blood pressure
15 and blood flow, whereas the neuronal enzyme (ncNOS) serves as a neurotransmitter and appears to be involved in the regulation of various biological functions such as cerebral ischaemia. Inducible NOS has been particularly implicated in the pathogenesis of inflammatory diseases. Regulation of these enzymes should therefore offer considerable potential in the treatment of a wide variety of disease states (J. E. Macdonald, Ann. Rep.
20 Med. Chem., 1996, 31, 221 - 230).


WO 97/14686 discloses, amongst other compounds, aminospiriperidine quinazoline derivatives of the following formula:



wherein R^3 represents various substituents, for use as pharmaceuticals. The treatment or prophylaxis of inflammatory conditions is disclosed as a particular pharmaceutical use.

5 Disclosure of the invention

According to the present invention, there is provided a compound of formula (I)



wherein:

- 10 R^1 and R^2 independently represent hydrogen, C1 to 6 alkyl, C2 to 6 alkenyl, C2 to 6 alkynyl, C1 to 6 alkoxy, C1 to 6 alkylthio, halogen, hydroxy, trifluoromethyl or amino;

- R^3 represents one or more substituents independently selected from hydrogen, C1 to 6 alkyl, C2 to 6 alkenyl, C2 to 6 alkynyl, C1 to 6 alkoxy, C1 to 6 alkylthio, halogen, hydroxy, trifluoromethyl, amino, cyano, nitro, trifluoromethoxy, methanesulphonyl, sulphonyl, $-NR^4R^5$, $-COOR^6$, $-CONR^7R^8$, benzyloxy, phenyl, or a 5-membered heterocyclic aromatic ring containing 1 to 3 heteroatoms which may be the same or different and are selected from O, N and S, which phenyl or 5-membered heterocyclic aromatic ring is optionally substituted, the optional substituents being C1 to 6 alkyl, halogen, cyano, nitro, hydroxy, C1 to 6 alkoxy, trifluoromethyl and trifluoromethoxy;
- 15
- 20

R⁴, R⁵ and R⁶ independently represent hydrogen or C1 to 6 alkyl;

R⁷ and R⁸ independently represent hydrogen, C1 to 6 alkyl or phenyl, which phenyl is
optionally substituted by one or more groups independently selected from C1 to 6 alkyl,
halogen, cyano, nitro, hydroxy, C1 to 6 alkoxy, trifluoromethyl and trifluoromethoxy;

X represents $-(CH_2)_n-$, wherein n represents zero or 1; and

A represents an aromatic carbocyclic ring or a 5- or 6-membered heterocyclic aromatic ring
containing 1 to 3 heteroatoms which may be the same or different and are selected from O,
N and S;

or a pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof.

The compounds of formula (I) and their pharmaceutically acceptable salts, enantiomers,
racemates and tautomers have the advantage that they are potent inhibitors of the inducible
isoform of the enzyme nitric oxide synthase (iNOS) present in macrophages.

The invention further provides a process for the preparation of such compounds or a
pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof.

According to the invention there is also provided a compound of formula (I), or a
pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof, for use as a
medicament.

Another aspect of the invention provides the use of a compound of formula (I) or a
pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof, in the
manufacture of a medicament, for the treatment or prophylaxis of diseases or conditions in
which inhibition of nitric oxide synthase activity is beneficial.

A more particular aspect of the invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof, in the manufacture of a medicament, for the treatment or prophylaxis of inflammatory disease.

5 According to the invention, there is also provided a method of treating, or reducing the risk of, diseases or conditions in which inhibition of nitric oxide synthase activity is beneficial which comprises administering to a person suffering from or at risk of, said disease or condition, a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof.

10

More particularly, there is also provided a method of treating, or reducing the risk of, inflammatory disease in a person suffering from or at risk of, said disease, wherein the method comprises administering to the person a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt, enantiomer, racemate or
15 tautomer thereof.

The compounds of the present invention may also be used advantageously in combination with a second pharmaceutically active substance, particularly in combination with a selective inhibitor of the inducible isoform of cyclooxygenase (COX-2). Thus, in a further
20 aspect of the invention there is provided the use of a compound of formula (I) or a pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof, in combination with a COX-2 inhibitor for the treatment of inflammation, inflammatory disease and inflammatory related disorders. And there is also provided a method of treating, or reducing the risk of, inflammation, inflammatory disease and inflammatory related
25 disorders in a person suffering from or at risk of, said disease or condition, wherein the method comprises administering to the person a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof in combination with a COX-2 inhibitor.

30 Preferably, A in formula (I) represents a benzo ring.

Preferably, R^1 in formula (I) represents hydrogen. Alternatively, it is preferred that R^1 in formula (I) represents fluoro.

5 Preferably, R^2 in formula (I) represents fluoro.

In another preferred embodiment, n in formula (I) has a value of zero.

Particular compounds of the invention include:

- 10 (R^*R^*)-4'-amino-5'-fluoro-1,3,4,10a-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)quinazoline]-6-one;
(R^*S^*)-4'-amino-5'-fluoro-1,3,4,10a-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)quinazoline]-6-one;
(R^*R^*)-4-amino-5-fluoro-3b',4',6',7'-tetrahydrospiro[quinazoline-2,5'(2H,9'H)-thieno[3,2-
15 a]indolizine]-9'-one;
(R^*S^*)-4-amino-5-fluoro-3b',4',6',7'-tetrahydrospiro[quinazoline-2,5'(2H,9'H)-thieno[3,2-a]indolizine]-9'-one;
(R^*R^*)-4-amino-5-fluoro-6',7',9',9a'-tetrahydrospiro[quinazoline-2,8'(2H,4'H)-thieno[2,3-a]indolizine]-4'-one;
20 (R^*S^*)-4-amino-5-fluoro-6',7',9',9a'-tetrahydrospiro[quinazoline-2,8'(2H,4'H)-thieno[2,3-a]indolizine]-4'-one;
(R^*R^*)-4'-amino-7-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
(R^*S^*)-4'-amino-7-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-
25 2,2'(1'H)-quinazoline]-6-one;
(R^*R^*)-4'-amino-8-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
(R^*S^*)-4'-amino-8-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;

- (*R***R*)-4'-amino-9-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R***S**)-4'-amino-9-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- 5 (*R***R**)-4'-amino-10-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R***S**)-4'-amino-10-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R***R**)-4'-amino-7,8-dichloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-10 2,2'(1'H)-quinazoline]-6-one;
- (*R***S**)-4'-amino-7,8-dichloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R***R**)-4'-amino-7-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- 15 (*R***S**)-4'-amino-7-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R***R**)-4'-amino-8-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R***S**)-4'-amino-8-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-20 2,2'(1'H)-quinazoline]-6-one;
- (*R***R**)-4'-amino-8-methoxy-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R***S**)-4'-amino-8-methoxy-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- 25 (*R***R**)-4'-amino-8-cyano-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R***S**)-4'-amino-8-cyano-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R***R**)-4'-amino-8-cyano-5',8'-difluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-30 2,2'(1'H)-quinazoline]-6-one;

(*R***S**)-4'-amino-8-cyano-5',8'-difluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;

(*R***R**)-4'-amino-5'-fluoro-7,8,10,10a-tetrahydrospiro[pyrido[3,4-a]indolizine-9,2'(1'H)-quinazoline]-5-one;

5 (*R***S**)-4'-amino-5'-fluoro-7,8,10,10a-tetrahydrospiro[pyrido[3,4-a]indolizine-9,2'(1'H)-quinazoline]-5-one;

(*R***R**)-4'-amino-5'-fluoro-6b,7,9,10-tetrahydrospiro[naphtho[2,1-a]indolizine-8,2'(1'H)-quinazoline]-12-one;

10 (*R***S**)-4'-amino-5'-fluoro-6b,7,9,10-tetrahydrospiro[naphtho[2,1-a]indolizine-8,2'(1'H)-quinazoline]-12-one;

and pharmaceutically acceptable salts, enantiomers, racemates or tautomers thereof.

Unless otherwise indicated, the term "C1 to 6 alkyl " referred to herein denotes a straight or branched chain alkyl group having from 1 to 6 carbon atoms or a cyclic alkyl group having
15 from 3 to 6 carbon atoms. Examples of such groups include methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, t-butyl, cyclopentyl and cyclohexyl.


Unless otherwise indicated, the term "C2 to 6 alkenyl " referred to herein denotes a straight or branched chain alkyl group having from 2 to 6 carbon atoms and including one double
20 bond or a cyclic alkyl group having from 3 to 6 carbon atoms and including one double bond. Examples of such groups include ethenyl, 1- and 2-propenyl, 2-methyl-2-propenyl, 2-butenyl, cyclopentenyl and cyclohexenyl.

Unless otherwise indicated, the term " C2 to 6 alkynyl " referred to herein denotes a
25 straight or branched chain alkyl group having from 2 to 6 carbon atoms and including one triple bond. Examples of such groups include ethynyl, 1- and 2-propynyl and 2-butylnyl.


Unless otherwise indicated, the term "C1 to 6 alkoxy " referred to herein denotes a straight or branched chain alkoxy group having from 1 to 6 carbon atoms. Examples of such groups
30 include methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, i-butoxy, s-butoxy and t-butoxy.

Other groups, for example, alkylthio, are to be interpreted similarly.

The process mentioned above, for the preparation of compounds of the invention, or a
5 pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof comprises
reaction of a compound of formula (II)



wherein R¹ and R² are as defined above,
10 with a compound of formula (III) or a protected derivative thereof



wherein R³, A and X are as defined above.

15 The reaction may be carried out in a polar solvent, for example, methanol, ethanol, acetonitrile, dimethylformamide or dimethylsulphoxide at a suitable temperature, generally between 20 °C and the boiling point of the solvent, or without solvent at a temperature generally between 20 °C and 200 °C. We have found that it is sometimes convenient to use
20 the compounds of formula (III) in a protected form, for example as an acetal such as the diethoxy acetal. The process is then preferably carried out in the presence of an acid catalyst. The required acetals may be formed by reacting an unprotected compound of formula (III) with an alcohol such as ethanol using methods that are well known in the art.

The present invention includes compounds of formula (I) in the form of salts, in particular acid addition salts. Suitable salts include those formed with both organic and inorganic acids. Such acid addition salts will normally be pharmaceutically acceptable although salts of non-pharmaceutically acceptable acids may be of utility in the preparation and
5 purification of the compound in question. Thus, preferred salts include those formed from hydrochloric, hydrobromic, sulphuric, phosphoric, citric, tartaric, lactic, pyruvic, acetic, succinic, fumaric, maleic, methanesulphonic and benzenesulphonic acids.


Salts of compounds of formula (I) may be formed by reacting the free base, or a salt,
10 enantiomer, racemate or tautomer thereof, with one or more equivalents of the appropriate acid. The reaction may be carried out in a solvent or medium in which the salt is insoluble or in a solvent in which the salt is soluble, for example, water, dioxane, ethanol, tetrahydrofuran or diethyl ether, or a mixture of solvents, which may be removed *in vacuo* or by freeze drying. The reaction may also be a metathetical process or it may be carried out on an ion
15 exchange resin.

Novel intermediates of formulae (II) and (III) form another aspect of the invention.

Compounds of formula (II) may be prepared using methods such as those described in WO
20 97/14686.

The preparation of compounds of formula (III) is either known *per se* or may be achieved using methods well known in the art. For example, compounds of formula (III) may be obtained by cyclisation of a compound of formula (IV)


25



wherein R^3 , A and X are as defined above and M is an alkali or alkali earth metal.

Examples of such metals M include lithium, sodium, potassium or magnesium.

Compounds of formula (IV) may be formed *in situ* by treatment of a compound of formula
 5 (IV')




wherein Hal represents a halogen atom, particularly bromine or iodine,


10 with the metal M or an organometallic derivative or salt thereof, using procedures well known to the art. The reactions are best carried out in an aprotic solvent such as ether, tetrahydrofuran or diglyme at temperatures between -100 °C and 30 °C.

Compounds of formula (IV') may be prepared by reaction of the compound of formula (V)

15



with an acid derivative of formula (VI)



20

wherein R^3 , A, X and Hal are as defined above, and L is a leaving group. The reaction may be performed in an organic solvent, for example ethanol, dichloromethane or

dimethylformamide at a temperature between 0 °C and the boiling point of the solvent. The reaction may be catalysed by the addition of a base; bases that may be used include organic amines (for example, triethylamine or pyridine) and alkali metal hydroxides, alkoxides or hydrides. Suitable leaving groups include halogen (especially chlorine) and imidazole.

- 5 These acid derivatives may be prepared from the corresponding acid (compounds of formula (VI) wherein L is OH) using methods well known in the art.

Intermediate compounds may be used in protected form. Protecting groups and details of processes for their removal may be found by reference to the standard text "Protecting groups
10 in Organic Synthesis", 2nd Edition (1991) by Greene and Wuts.

The compounds of the invention and intermediates thereto may be isolated from their reaction mixtures and, if necessary further purified, by using standard techniques.

- 15 The compounds of formula I may exist in enantiomeric forms. Therefore, all enantiomers, diastereomers, racemates and mixtures thereof are included within the scope of the invention. The various optical isomers may be isolated by separation of a racemic mixture of the compounds using conventional techniques, for example, fractional crystallisation, or HPLC.

- 20 Intermediate compounds may also exist in enantiomeric forms and may be used as purified enantiomers, diastereomers, racemates or mixtures.

The compounds of formula (I) may exist in alternative tautomeric forms. Compounds of formula (I) are provided in another tautomeric form or as a mixture thereof.

25

The compounds of formula (I), and their pharmaceutically acceptable salts, enantiomers, racemates and tautomers, are useful because they possess pharmacological activity in animals. In particular, the compounds are active as inhibitors of the enzyme nitric oxide synthase. More particularly, they are inhibitors of the inducible isoform of the enzyme nitric

oxide synthase present in macrophages and as such are predicted to be useful in therapy, for example, as anti-inflammatory agents.

The compounds and their pharmaceutically acceptable salts, enantiomers, racemates and
5 tautomers are indicated for use in the treatment or prophylaxis of diseases or conditions in which synthesis or oversynthesis of nitric oxide synthase forms a contributory part. In particular, the compounds are indicated for use in the treatment of inflammatory conditions in mammals including man.

10 Conditions that may be specifically mentioned are:

osteoarthritis, rheumatoid arthritis, rheumatoid spondylitis, gouty arthritis and other arthritic conditions, inflamed joints;

15 eczema, psoriasis, dermatitis or other inflammatory skin conditions such as sunburn;

inflammatory eye conditions including uveitis and conjunctivitis;

lung disorders in which inflammation is involved, for example, asthma, bronchitis, pigeon
20 fancier's disease, farmer's lung, acute respiratory distress syndrome;

bacteraemia, endotoxaemia (septic shock), aphthous ulcers, gingivitis, pyresis, pain and pancreatitis;

25 conditions of the gastrointestinal tract including Crohn's disease, atrophic gastritis, gastritis varialoforme, ulcerative colitis, coeliac disease, regional ileitis, peptic ulceration, irritable bowel syndrome, damage to the gastrointestinal tract resulting from infections by, for example, *Helicobacter pylori*, or from treatments with non-steroidal anti-inflammatory drugs;

30 and other conditions associated with inflammation.

The compounds will also be useful in the treatment and alleviation of acute or persistent inflammatory or neuropathic pain or pain of a central origin.

5 The compounds of formula (I) and their pharmaceutically acceptable salts, enantiomers, racemates and tautomers may also be useful in the treatment or prophylaxis of diseases or conditions in addition to those mentioned above. For example, the compounds may be useful in the treatment of atherosclerosis, cystic fibrosis, hypotension associated with septic and/or toxic shock, in the treatment of dysfunction of the immune system, as an adjuvant to short-
10 term immunosuppression in organ transplant therapy, in the treatment of vascular complications associated with diabetes and in cotherapy with cytokines, for example TNF or interleukins.

The compounds of formula (I) may also show inhibitory activity against the neuronal isoform
15 of nitric oxide synthase. Thus they may also be useful in the treatment of hypoxia, for example in cases of cardiac arrest and stroke, neurodegenerative disorders including nerve degeneration and/or nerve necrosis in disorders such as hypoxia, hypoglycaemia, epilepsy, and in external wounds (such as spinal cord and head injury), hyperbaric oxygen convulsions and toxicity, dementia, for example pre-senile dementia, Alzheimer's disease and AIDS-
20 related dementia, Sydenham's chorea, Parkinson's disease, Tourette's Syndrome, Huntington's disease, Amyotrophic Lateral Sclerosis, Korsakoff's disease, imbecility relating to a cerebral vessel disorder, sleeping disorders, schizophrenia, depression, autism, seasonal affective disorder, jet-lag, depression or other symptoms associated with Premenstrual Syndrome (PMS), anxiety and septic shock. Compounds of formula (I) may also be expected to show
25 activity in the prevention and reversal of tolerance to opiates and diazepines, treatment of drug addiction, treatment of migraine and other vascular headaches, neurogenic inflammation, in the treatment of gastrointestinal motility disorders, cancer and in the induction of labour.

30 For the above mentioned therapeutic indications, the dosage administered will, of course, vary with the compound employed, the mode of administration and the treatment desired.

However, in general, satisfactory results are obtained when the compounds are administered at a dosage of the solid form of between 1 mg and 2000 mg per day.

The compounds of formula (I), and pharmaceutically acceptable derivatives thereof, may be used on their own, or in the form of appropriate pharmaceutical compositions in which the compound or derivative is in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier. Administration may be by, but is not limited to, enteral (including oral, sublingual or rectal), intranasal, intravenous, topical or other parenteral routes. Conventional procedures for the selection and preparation of suitable pharmaceutical formulations are described in, for example, "Pharmaceuticals - The Science of Dosage Form Designs", M. E. Aulton, Churchill Livingstone, 1988. The pharmaceutical composition preferably comprises less than 80% and more preferably less than 50% of a compound of formula (I), or a pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof.

There is also provided a process for the preparation of such a pharmaceutical composition which comprises mixing the ingredients.

The compounds of formula (I), and pharmaceutically acceptable derivatives thereof, may also be advantageously used in combination with a COX-2 inhibitor. Particularly preferred COX-2 inhibitors are Celecoxib and MK-966. The NOS inhibitor and the COX-2 inhibitor may either be formulated together within the same pharmaceutical composition for administration in a single dosage unit, or each component may be individually formulated such that separate dosages may be administered either simultaneously or sequentially.

The invention is illustrated, but in no way limited, by the following examples:

Preparation 1

2,3-Dichloro-6-iodobenzoic acid chloride

a) 2,3-Dichloro-6-iodobenzoic acid

n-Butyllithium (1.45M solution in hexane, 19.4 mmol, 13.4 ml) was added dropwise to a stirred solution of diisopropylamine (21.4 mmol, 3.0 ml) in THF (90 ml) at 0 °C. After 30 minutes, the solution was cooled to -78 °C and a solution of 1,2-dichloro-4-iodobenzene
5 (5.30 g, 19.4 mmol) in THF (10 ml) was added dropwise. After 10 minutes, solid carbon dioxide was added cautiously and the mixture allowed to warm to room temperature over 16 h. The mixture was diluted with water and extracted twice with ethyl acetate. The aqueous layer was acidified with dilute aqueous hydrochloric acid and then extracted twice with ethyl acetate. The extracts were dried over sodium sulphate and evaporated to give
10 the title compound as a yellow solid (3.59 g). 300 MHz ¹H NMR (d₆-DMSO) 11.5 (1H, br.s), 7.82 (1H, d, J 8.4 Hz), 7.44 (1H, d, J 8.4 Hz).

b) 2,3-Dichloro-6-iodobenzoic acid chloride

DMF (one drop) was added to a stirred solution of 2,3-dichloro-6-iodobenzoic acid (1.53 g, 4.83 mmol) and oxalyl chloride (2 ml) in ethyl acetate (20 ml). After 2 hours, the solution
15 was evaporated and the product was used immediately.

Preparation 23-Cyano-6-iodobenzoic acid chloride20 a) 2-Amino-5-cyanobenzoic acid methyl ester

2-Amino-5-bromobenzoic acid methyl ester (4.6 g, 20 mmol) and copper (I) cyanide (3.6 g, 40 mmol) were heated at 200 °C in N-methylpyrrolidinone (20 ml) for 4 h. After cooling to room temperature, the solution was diluted with water, extracted with ethyl acetate, the aqueous extract acidified with 2N hydrochloric acid and filtered. The organic layer was
25 separated, dried over magnesium sulphate and evaporated. The residue was purified by flash column chromatography on silica eluting with 50% ethyl acetate in hexane to yield the title compound as a colourless oil (1.0 g). MS (+EI) ^{m/z} 176 [M⁺].

b) 2-Iodo-5-cyanobenzoic acid methyl ester

Sodium nitrite (0.8 g, 11.4 mmol) in water (10 ml) was added dropwise over 10 minutes to
30 a stirred solution of 2-amino-5-cyanobenzoic acid methyl ester (2.0 g, 11.4 mmol) in water

(30 ml) and conc. sulphuric acid (15 ml) at 0 °C. After a further 15 minutes at 5 °C, a solution of potassium iodide (3.5 g, 21 mmol) in water (30 ml) was quickly added and the mixture allowed to warm to room temperature. 10% Aqueous sodium thiosulphate was added, the mixture extracted with ethyl acetate, the extract dried over magnesium sulphate and evaporated. The residue was purified by flash column chromatography on silica eluting with 20% ethyl acetate in hexane to yield the title compound as a colourless oil (2.8 g). MS (+EI) m/z 287 $[M]^+$.

c) 2-Iodo-5-cyanobenzoic acid

A solution of sodium hydroxide (0.44 g, 11 mmol) in water (10 ml) was added to 2-iodo-5-cyanobenzoic acid methyl ester (2.7 g, 9.4 mmol) in methanol. The mixture was stirred for 20 h, then most of the solvent was evaporated and the residue acidified with 2N hydrochloric acid. The resulting precipitate was filtered off and dried to give a white solid (1.7 g). MS (-CI) m/z 272 $[M-H]^+$.

d) 2-Iodo-5-cyanobenzoic acid chloride

This was prepared by the method of Preparation 1(b) and was used directly.

Other acid chlorides are either known *per se* or were prepared from the known acids by the method of Preparation 1(b).

Preparation 3

1-(2-Bromo-6-chlorobenzoyl)-2,3-dihydro-4-pyridinone

A solution of 2-bromo-6-chlorobenzoic acid chloride (0.5 mmol) in dichloromethane (5 ml) was added dropwise to a solution of 2,3-dihydro-4-pyridinone (5.1 mmol, 495 mg) and triethylamine (1.2 ml) in dichloromethane (20 ml). After stirring for 30 minutes, the solution was diluted with water, separated, the aqueous extract further extracted with dichloromethane, the combined organic extracts combined and washed with brine, dried over sodium sulphate and evaporated. The residue was purified by flash column chromatography on silica eluting with 30% ethyl acetate in hexane to yield the title compound as a pale yellow oil (629 mg). MS (+CI) m/z 316 $[M+H]^+$.

The following intermediates were similarly prepared from 2,3-dihydro-4-pyridinone and the appropriate acid chloride:

- 5 (i) 1-(2-Bromo-5-methylbenzoyl)-2,3-dihydro-4-pyridinone: colourless oil, MS (+EI) m/z 293/295 [M^+].
- (ii) 1-(3-Bromo-4-pyridinoyl)-2,3-dihydro-4-pyridinone: white solid, MS (+EI) m/z 280/282 [M^+].
- 10 (iii) 1-(2,3-Dichloro-6-iodobenzoyl)-2,3-dihydro-4-pyridinone: white solid, MS (+EI) m/z 395/397 [M^+].
- (iv) 1-(5-Cyano-2-iodobenzoyl)-2,3-dihydro-4-pyridinone: colourless oil, used without
15 purification.
- (v) 1-(2-Methyl-6-iodobenzoyl)-2,3-dihydro-4-pyridinone : MS (+EI) m/z 341 [M^+].
- (vi) 1-(2-Bromo-3-chlorobenzoyl)-2,3-dihydro-4-pyridinone: MS (+EI) m/z 313/315
20 [M^+].
- (vii) 1-(2-Bromo-4-chlorobenzoyl)-2,3-dihydro-4-pyridinone: MS (+EI) m/z 314 [M^+].
- (viii) 1-(2-Bromo-5-chlorobenzoyl)-2,3-dihydro-4-pyridinone: MS (+EI) m/z 314 [M^+].
- 25 (ix) 1-(2-Bromo-3-thienoyl)-2,3-dihydro-4-pyridinone: MS (-CI) m/z 284/286 [$M-H$] $^+$.

(x) 1-(2-Bromo-5-methoxybenzoyl)-2,3-dihydro-4-pyridinone; MS (+EI) m/z 309/311 $[M^+]$.

(xi) 1-(2-Iodo-1-naphthoyl)-2,3-dihydro-4-pyridinone: MS (-CI) m/z 376 $[M-H]^+$.

5

(xii) 1-(2-Iodo-3-thienoyl)-2,3-dihydro-4-pyridinone: colourless oil : 300 MHz 1H NMR (d_6 -DMSO) 7.55 (1H, d, J 5.4 Hz), 7.47-7.43 (1H, m), 7.00 (1H, d, J 5.4 Hz), 5.40 (1H, d, J 8.1 Hz), 4.18 (2H, t, J 7.2 Hz), 2.68 (2H, t, J 7.2 Hz).

10

Preparation 4

(±)-7-Chloro-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione

A mixture of 1-(2-bromo-6-chlorobenzoyl)-2,3-dihydro-4-pyridinone (Preparation 3) (300 mg, 0.954 mmol), triethylamine (0.39 mmol, 0.54 ml), formic acid (0.05 ml, 1.3 mmol) and di(triphenylphosphine) diacetoxypalladium (47 mg, 0.063 mmol) in DMF (3 ml) was heated to 80 °C for 2 h. The cooled solution was filtered through celite, diluted with water, extracted twice with ethyl acetate, the extracts dried over sodium sulphate and evaporated. The residue was purified by flash column chromatography on silica eluting with 50% ethyl acetate in hexane to yield the title compound as a colourless oil (70 mg). MS (+CI) m/z 235/237 $[M+H]^+$.

20

The compounds of Preparations 5 to 7 were synthesised using the method of Preparation 4:

Preparation 5

25

(±)-7,8-Dichloro-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione

From 1-(2,3-dichloro-6-iodobenzoyl)-2,3-dihydro-4-pyridinone. White solid. MS (+EI) m/z 269/271 $[M^+]$.

Preparation 6(±)-8-Chloro-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione

- 5 From 1-(2-bromo-5-chlorobenzoyl)-2,3-dihydro-4-pyridinone. Colourless oil. MS (+EI) m/z 235/237 [M^+].

Preparation 7

10 (±)-3b,4,6,7-tetrahydrothieno[3,2-a]indolizine-5,9-dione

From 1-(3-bromo-2-thienoyl)-2,3-dihydro-4-pyridinone. Colourless oil. MS (+CI) m/z 208 [$M+H$]⁺.

Preparation 8

15

(±)-8-Methyl-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione

- A mixture of 1-(2-bromo-5-methylbenzoyl)-2,3-dihydro-4-pyridinone (1.19 g, 4.03 mmol), tri-n-butyltin hydride (1.14 ml, 4.24 mmol) and azobis(isobutyronitrile) (AIBN, 66 mg, 10 mol%) in toluene (40 ml) was heated at 95 °C for 2 h, cooled, evaporated and
20 purified by flash column chromatography on silica eluting with 30% ethyl acetate in hexane to yield the title compound as a pale yellow oil (276 mg), MS (+EI) m/z 215 [M^+].

The compounds of Preparations 9 to 12 were prepared using the method of Preparation 8:

25

Preparation 9(±)-8-Cyano-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione:

From 1-(5-cyano-2-iodobenzoyl)-2,3-dihydro-4-pyridinone. MS (-CI) m/z 226 [M^+].

Preparation 10(±)-7,8,10,10a-Tetrahydropyrido[3,4-a]indolizine-5,9-dione

From 1-(3-bromo-4-pyridinoyl)-2,3-dihydro-4-pyridinone. MS (+EI) m/z 202 $[M^+]$.

5

Preparation 11(±)-7-Methyl-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione

From 1-(2-Methyl-6-iodobenzoyl)-2,3-dihydro-4-pyridinone. MS (+EI) m/z 215 $[M^+]$.

10

Preparation 12(±)-6b,7,9,10-tetrahydronaphtho[2,1-a]indolizine-8,12-dione

From 1-(2-iodo-1-naphthoyl)-2,3-dihydro-4-pyridinone. MS (+CI) m/z 252 $[M+H]^+$.

15

Preparation 13(±)-8-Methoxy-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione

Butyl lithium (3.1 ml, 0.9 M in hexanes, 2.8 mmol) was added dropwise to a solution of
20 1-(2-bromo-5-methoxybenzoyl)-2,3-dihydro-4-pyridinone (0.75 g, 2.4 mmol) in THF (20 ml) at -78 °C. After stirring for 2 h, the reaction was quenched with water, extracted with ethyl acetate, the extract dried over magnesium sulphate and evaporated. The residue was purified by flash column chromatography on silica eluting with 50% ethyl acetate in hexane to yield the title compound as a colourless oil (240 mg). MS (+EI) m/z 231 $[M^+]$.

25

The compounds of Preparations 14 to 16 were prepared using the method of Preparation 13:

Preparation 1410-Chloro-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione

From 1-(2-bromo-3-chlorobenzoyl)-2,3-dihydro-4-pyridinone: colourless oil. MS (+EI) m/z
5 235/237 [M^+].

Preparation 159-Chloro-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione


10 From 1-(2-bromo-4-chlorobenzoyl)-2,3-dihydro-4-pyridinone: colourless oil. MS (+EI) m/z
235/237 [M^+].

Preparation 166,7,9,9a-Tetrahydrothieno[2,3-a]indolizine-4,8-dione

From 1-(2-bromo-3-thienoyl)-2,3-dihydro-4-pyridinone. MS (+CI) m/z 208 [$M+H$]⁺.

Example 1

20 (±)-(R*R*)-4'-Amino-5'-fluoro-1,3,4,10a-tetrahydrospiro[pyrido[2,1-a]isoindole-
2,2'(1'H)quinazoline]-6-one




A solution of 2-amino-6-fluorobenzamidine (0.13 g, 0.61 mmol) and 1,3,4,10a-tetrahydro[pyrido[2,1-a]isoindole-2,6-dione (0.12 g, 0.59 mmol) (J.C.S. Perkin Trans. 1,

1984, 2477; Tetrahedron, 1993, 2239) in ethanol (15 ml) was stirred at reflux for 18 h. The mixture was cooled and filtered to give a yellow solid (0.1 g), which was purified by flash column chromatography on neutral alumina, eluting with dichloromethane/methanol (99:1 to 95:5) as eluent to afford a single diastereoisomer of the title compound (containing the
5 RR and SS enantiomers), as a yellow solid (0.05g). M.p. 207-209 °C.

Example 2

10 (±)-(R*S*)-4'-Amino-5'-fluoro-1,3,4,10a-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)quinazoline]-6-one




The filtrate obtained after the precipitation of the product of Example 1 was concentrated to give a brown oil (0.19 g). Purification by flash column chromatography on neutral
15 alumina, eluting with dichloromethane/methanol (99:1 to 95:5) as eluent gave a second diastereoisomer of the title compound (containing the RS and SR enantiomers), as a yellow solid (0.03g). M.p.184-186 °C.

Example 3

20

(±)-(R*R*)-4-Amino-5-fluoro-3b',4',6',7'-tetrahydrospiro[quinazoline-2,5'(2H,9'H)-thieno[3,2-a]indolizine]-9'-one




A solution of 2-amino-6-fluorobenzamidine dihydrochloride (100 mg., 44 mmol) and 3b,4,6,7-tetrahydrothieno[3,2-a]indolizine-5,9-dione (Preparation 7, 80 mg, 39 mmol) in ethanol (10 ml) was heated at reflux for 5 h. The solvent was evaporated and the residue
 5 separated by chromatography on silica using dichloromethane-methanol mixtures as eluent. The first eluted fraction provided the title compound (63 mg, 43%) as a pale yellow solid, m.p. 193-195 °C. MS (+CI) m/z 343 $[M+H]^+$.

Example 4

10

(±)-(R*S*)-4-Amino-5-fluoro-3b',4',6',7'-tetrahydrospiro[quinazoline-2,5'(2H,9'H)-thieno[3,2-a]indolizine]-9'-one




The second eluted fraction from the chromatography described in Example 3 afforded the
 15 title compound (33 mg, 22%), m.p. 218-220 °C. MS (+CI) m/z 343 $[M+H]^+$.

The compounds of Examples 5 to 30 were made by the method of Examples 3 and 4 using the intermediates stated.

20 The compounds of Examples 5 and 6 were obtained from 2-amino-6-fluorobenzamidine dihydrochloride and 6,7,9,9a-tetrahydrothieno[2,3-a]indolizine-4,8-dione (Preparation 16).

Example 5

5 (±)-(R*R*)-4-Amino-5-fluoro-6',7',9',9a'-tetrahydrospiro[quinazoline-2,8'(2H,4'H)-thieno[2,3-a]indolizine]-4'-one




Obtained as the hydrochloride salt, m.p. >275 °C. MS (+CI) m/z 343 $[M+H]^+$.

Example 6

10

(±)-(R*S*)-4-Amino-5-fluoro-6',7',9',9a'-tetrahydrospiro[quinazoline-2,8'(2H,4'H)-thieno[2,3-a]indolizine]-4'-one




Obtained as the hydrochloride salt, m.p. 225-230 °C. MS (+CI) m/z 343 $[M+H]^+$.

15

The compounds of Examples 7 and 8 were obtained from 2-amino-6-fluorobenzamidine dihydrochloride and 7-chloro-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione (Preparation 4).

Example 7


(±)-(R*R*)-4'-Amino-7-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one hydrochloride



5 MS (+Cl) m/z 371/373 $[M+H]^+$; 300 MHz 1H NMR (d_6 -DMSO) 7.62 (1H, d, J 7.5 Hz), 7.54 (1H, t, J 7.5 Hz), 7.45 (1H, d, J 7.2 Hz), 7.13 (1H, dt, J 6.6, 6.6, 8.1 Hz), 6.59 (1H, s), 6.40 (1H, d, J 7.2 Hz), 6.37-6.33 (1H, m), 5.85-5.8 (2H, br.s), 4.79 (1H, dd, J 3.3, 11.4 Hz), 4.12 (1H, dd, J 3.9, 13.8 Hz), 3.40-3.35 (1H, m), 2.60-2.55 (1H, m), 1.91 (1H, d, J 12 Hz),
10 1.37 (1H, dt, J 5.7, 12.9, 12.9 Hz), 1.04 (1H, t, J 11.7 Hz).

Example 8

15 (±)-(R*S*)-4'-Amino-7-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one hydrochloride




MS (+Cl) m/z 371/373 $[M+H]^+$; 300 MHz 1H NMR (d_6 -DMSO) 7.61-7.46 (4H, m), 7.27-7.20 (2H, m), 6.74 (1H, d, J 8.1 Hz), 6.44 (1H, dd, J 8.4, 12.3 Hz), 4.75 (1H, dd, J 3.6, 3.9 Hz), 3.32-3.19 (1H, m), 2.0-1.95 (1H, m), 1.5-1.4 (1H, m), 1.3-1.1 (1H, m).

The compounds of Examples 9 and 10 were obtained from 2-amino-6-fluorobenzamidine dihydrochloride and 8-chloro-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione (Preparation 6).

5

Example 9

(±)-(R*R*)-4'-Amino-8-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




10 Obtained as the hydrochloride salt, m.p. 308 °C (dec.).

Example 10

(±)-(R*S*)-4'-Amino-8-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one

15



Obtained as the trifluoroacetate salt (amorphous) after purification by reverse-phase HPLC.


MS (+Cl) m/z 371 $[M+H]^+$; MS (-Cl) m/z 369 $[M-H]^+$.

The compounds of Examples 11 and 12 were obtained from 2-amino-6-fluorobenzamidine dihydrochloride and 9-chloro-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione (Preparation 15).

5

Example 11


(±)-(R*R*)-4'-Amino-9-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one



10 Obtained as the hydrochloride salt, m.p. 225 °C (dec).

Example 12

15 (±)-(R*S*)-4'-Amino-9-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




Obtained as the hydrochloride salt, m.p. 210 °C (dec).

20 The compounds of Examples 13 and 14 were obtained from 2-amino-6-fluorobenzamidine dihydrochloride and 10-chloro-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione (Preparation 14).

Example 13


(±)-(R*R*)-4'-Amino-10-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one



Obtained as the hydrochloride salt. MS (+Cl) m/z 371 $[M+H]^+$.

Example 14

(±)-(R*S*)-4'-Amino-10-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




Obtained as the hydrochloride salt. MS (+Cl) m/z 371 $[M+H]^+$.

The compounds of Examples 15 and 16 were prepared from 2-amino-6-fluorobenzamidine dihydrochloride and 7,8-dichloro-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione (Preparation 5).

Example 15


(±)-(R*R*)-4'-Amino-7,8-dichloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one



Obtained as the hydrochloride salt, m.p. 258-260 °C.

Example 16

(±)-(R*S*)-4'-Amino-7,8-dichloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




Obtained as the hydrochloride salt, m.p. 259-261 °C.

- 15 The compounds of Examples 17 and 18 were obtained from 2-amino-6-fluorobenzamidine dihydrochloride and 7-methyl-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione (Preparation 11).

Example 17

(±)-(R*R*)-4'-Amino-7-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




5

Obtained as the hydrochloride salt, m.p. 201-204 °C.

Example 18

10 (±)-(R*S*)-4'-Amino-7-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




Obtained as the hydrochloride salt, m.p. 215-218 °C.

15 The compounds of Examples 19 and 20 were obtained from 2-amino-6-fluorobenzamidine dihydrochloride and 8-methyl-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione (Preparation 8).

Example 19


(±)-(R*R*)-4'-Amino-8-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one



Obtained as the hydrochloride salt, m.p. 238-239 °C.

Example 20

(±)-(R*S*)-4'-Amino-8-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




Obtained as the hydrochloride salt, m.p. 222-224 °C.

- 15 The compounds of Examples 21 and 22 were prepared from 2-amino-6-fluorobenzamidine dihydrochloride and 8-methoxy-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione (Preparation 13).

Example 21

(±)-(R*R*)-4'-Amino-8-methoxy-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




5

Obtained as the hydrochloride salt. MS (+CI) m/z 367 $[M+H]^+$.

Example 22

10 (±)-(R*S*)-4'-Amino-8-methoxy-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




Obtained as the hydrochloride salt. MS (+CI) m/z 367 $[M+H]^+$.

15 The compounds of Examples 23 and 24 were obtained from 2-amino-6-fluorobenzamidine dihydrochloride and 8-cyano-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione (Preparation 9).

Example 23


(±)-(R*R*)-4'-Amino-8-cyano-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one



Obtained as the hydrochloride salt, m.p. 241 °C (dec).

Example 24

(±)-(R*S*)-4'-Amino-8-cyano-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




Obtained as the hydrochloride salt, m.p. 290 °C (dec).

- 15 The compounds of Examples 25 and 26 were obtained from from 2-amino-3,6-difluoro benzamidine hydrochloride and 8-cyano-1,3,4,10b-tetrahydropyrido[2,1-a]isoindole-2,6-dione (Preparation 9).

Example 25

(±)-(R*R*)-4'-Amino-8-cyano-5',8'-difluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




5

Obtained as the hydrochloride salt, m.p. 243 °C (dec).

Example 26

10 (±)-(R*S*)-4'-Amino-8-cyano-5',8'-difluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one




M.p. 180 °C (dec).

15 The compounds of Examples 27 and 28 were obtained from 2-amino-6-fluorobenzamidine dihydrochloride and 7,8,10,10a-tetrahydropyrido[3,4-a]indolizine-5,9-dione (Preparation 10).

Example 27


(±)-(R*R*)-4'-Amino-5'-fluoro-7,8,10,10a-tetrahydrospiro[pyrido[3,4-a]indolizine-9,2'(1'H)-quinazoline]-5-one



Obtained as the hydrochloride salt. MS (+CI) m/z 338 $[M+H]^+$; 300 MHz 1H NMR (d_6 -DMSO) 8.97 (1H, s), 8.72 (1H, d, J 5.1 Hz), 7.67 (1H, d, J 5 Hz), 7.20 (1H, q, J 7 Hz), 6.8 (1H, br.s), 6.46-6.38 (2H, m), 4.97 (1H, dd, J 3.6, 12 Hz), 4.18 (1H, dd, J 4.5 12.6 Hz), 3.45 (1H, dt, J 3.3, 12.9 Hz), 2.76-2.72 (1H, m), 1.97 (1H, d, J 12 Hz), 1.45 (1H, dt, J 5.4, 12.9 Hz), 1.10 (1H, t, J 12.0 Hz).

Example 28

(±)-(R*S*)-4'-Amino-5'-fluoro-7,8,10,10a-tetrahydrospiro[pyrido[3,4-a]indolizine-9,2'(1'H)-quinazoline]-5-one




MS (+CI) m/z 338 $[M+H]^+$; 300 MHz 1H NMR (d_6 -DMSO) 8.95 (1H, s), 8.74 (1H, d, J 4.8 Hz), 7.69 (1H, d, J 5.1 Hz), 7.44-7.25 (3H, m), 6.78 (1H, d, J 7.5 Hz), 6.52-6.46 (1H, m), 4.94 (1H, dd, J 3.6, 12.3 Hz), 4.20 (1H, dd, J 4.8, 13.8 Hz), 3.8-3.43 (1H, dd, obscured by

H₂O), 2.71 (1H, br.d, J 9 Hz), 1.98 (1H, d, J 13.2 Hz), 1.55 (1H, dt, J 5.1, 12.9 Hz), 1.26 (1H, t, J 12.9 Hz).

The compounds of Examples 29 and 30 were obtained from 2-amino-6-fluorobenzamidine dihydrochloride and 6b,7,9,10-tetrahydronaphtho[2,1-a]indolizine-8,12-dione (Preparation 12).

Example 29

10 (±)-(R*R*)-4'-Amino-5'-fluoro-6b,7,9,10-tetrahydrospiro[naphtho[2,1-a]indolizine-8,2'(1'H)-quinazoline]-12-one




Obtained as the hydrochloride salt, m.p. 208-210 °C.

15

Example 30

(±)-(R*S*)-4'-Amino-5'-fluoro-6b,7,9,10-tetrahydrospiro[naphtho[2,1-a]indolizine-8,2'(1'H)-quinazoline]-12-one



20 Obtained as the hydrochloride salt, m.p. 235-237 °C.

Screens

The pharmacological activity of compounds according to the invention was tested in the following screens.

Screen 1

The activity of compounds of formula (I), or a pharmaceutically acceptable salt, enantiomer or tautomer thereof, may be screened for nitric oxide synthetase inhibiting activity by a procedure based on that of Förstermann *et al.*, Eur. J. Pharm., 1992, **225**, 161-165. Nitric oxide synthase converts ^3H -L-arginine into ^3H -L-citrulline which can be separated by cation exchange chromatography and quantified by liquid scintillation counting.

Enzyme is prepared, after induction, from the cultured murine macrophage cell line J774A-1 (obtained from the laboratories of the Imperial Cancer Research Fund). J774A-1 cells are cultured in Dulbeccos Modified Eagles Medium (DMEM) supplemented with 10% foetal bovine serum, 4 mM L-glutamine and antibiotics (100 units/ml penicillin G, 100 mg/ml streptomycin & 0.25 mg/ml amphotericin B). Cells are routinely grown in 225 cm³ flasks containing 35 ml medium kept at 37 °C and in a humidified atmosphere containing 5% CO₂.

Nitric oxide synthase is produced by cells in response to interferon-g (IFNg) and lipopolysaccharide (LPS). The medium from confluent culture flasks is removed and replaced with 25 ml (per flask) of fresh medium containing 1 mg/ml LPS and 10 units/ml IFNg. After a period of 17-20 hours in culture, harvesting of cells is accomplished by scraping the cell sheet from the flask surface into the culture medium. Cells are collected by centrifugation (1000 g for 10 minutes) and lysate prepared by adding to the cell pellet a solution containing 50 mM Tris-HCl (pH 7.5 at 20 °C), 10% (v/v) glycerol, 0.1% (v/v) Triton-X-100, 0.1 mM dithiothreitol and a cocktail of protease inhibitors comprising leupeptin (2 mg/ml), soya bean trypsin inhibitor (10 mg/ml), aprotinin (5 mg/ml) and phenylmethylsulphonyl fluoride (50 mg/ml).

For the assay, 25 μ l of substrate cocktail (50 mM Tris-HCl (pH 7.5 at 20 °C), 400 μ M NADPH, 20 μ M flavin adenine dinucleotide, 20 μ M flavin mononucleotide, 4 μ M tetrahydrobiopterin, 12 μ M L-arginine and 0.025 mCi L-[3 H] arginine) is added to wells of a 96 well filter plate (0.45 μ M pore size) containing 25 μ l of a solution of test compound in 50 mM Tris-HCl. The reaction is started by adding 50 μ l of cell lysate (prepared as above) and after incubation for 1 hour at room temperature is terminated by addition of 50 μ l of an aqueous solution of 3 mM nitroarginine and 21 mM EDTA.

Labelled L-citrulline is separated from labelled L-arginine using Dowex AG-50W. 150 μ l of a 25% aqueous slurry of Dowex 50W (Na $^+$ form) is added to the assay after which the whole is filtered into 96 well plates. 75 μ l of filtrate is sampled and added to wells of 96 well plates containing solid scintillant. After allowing the samples to dry the L-citrulline is quantified by scintillation counting.

In a typical experiment basal activity is 300 dpm per 75 μ l sample which is increased to 1900 dpm in the reagent controls. Compound activity is expressed as IC $_{50}$ (the concentration of drug substance which gives 50% enzyme inhibition in the assay) and aminoguanidine, which gives an IC $_{50}$ (50% inhibitory concentration) of 10 μ M, is tested as a standard to verify the procedure. Compounds are tested at a range of concentrations and from the inhibitions obtained IC $_{50}$ values are calculated. Compounds that inhibit the enzyme by at least 25% at 100 μ M are classed as being active and are subjected to at least one retest.

In the above screen, the compounds of Examples 1 to 30 were tested and gave IC $_{50}$ values of less than 25 μ M indicating that they are expected to show useful therapeutic activity.

Screen 2

Compounds also show activity against the human form of induced nitric oxide synthase as can be demonstrated in the following assay.

Enzyme is prepared, after induction, from the cultured human colon adenocarcinoma cell line DLD1 (obtained from the European Collection of Animal Cell Culture - cell line number 90102540). DLD1 cells are cultured in RPMI 1640 medium supplemented with 10% foetal bovine serum, 4 mM L-glutamine and antibiotics (100 units/ml penicillin G, 100 µg/ml streptomycin and 0.25 µg/ml amphotericin B). Cells are routinely grown in 225 cm³ flasks containing 35 ml medium kept at 37 °C and in a humidified atmosphere containing 5% CO₂.

Nitric oxide synthase is produced by cells in response to interferon-γ (IFN-γ) and interleukin-1β (IL-1β). The medium from confluent flasks is removed and replaced with 25 ml (per flask) of fresh medium containing 250 units/ml IL-1β and 1000 units/ml IFN-γ. After a period of 17–20 hours in culture, harvesting of cells is accomplished by scraping the cell monolayer from the flask surface into the culture medium. Cells are collected by centrifugation (1000g for 10 minutes) and lysate prepared by adding to the cell pellet a solution containing 50 mM Tris-HCl (pH 7.5 at 20°C), 10% (v/v) glycerol, 0.1% (v/v) Triton-X100, 0.1 mM dithiothreitol and a cocktail of protease inhibitors including leupeptin (2 µg/ml), soya bean trypsin inhibitor (10 µg/ml), aprotonin (5 µg/ml) and phenylmethylsulphonyl fluoride (50 µg/ml).

For the assay, 25 µl of substrate cocktail (50 mM Tris-HCl (pH 7.5), 400 µM NADPH, 20 µM flavin adenine dinucleotide, 20 µM flavin mononucleotide and 4 µM tetrahydrobiopterin) is added to the wells of a 96-well plate. Test compounds are preincubated with enzyme by adding together with 40 µl of cell lysate (prepared as above) and incubating for 1 hour at 37 °C at the end of which period 10 µl of 30 µM L-arginine and 0.025 µCi of L-[³H]-arginine in 50 mM Tris-HCl is added to start the enzymatic reaction. Incubation is continued for a further 1 hour at 37 °C. The reaction is terminated by addition of 50 µl of an aqueous solution of 3 mM nitroarginine and 21 mM EDTA.

Labelled L-citrulline is separated from labelled L-arginine using Dowex AG-50W. 120 µl of a 25% aqueous slurry of Dowex 50W is added to 96 well filter plates (0.45 µm pore


size). To this is added 120 μ l of terminated assay mix. 75 μ l of filtrate is sampled and added to the wells of 96 well plates containing solid scintillant. After allowing the samples to dry the L-citrulline is quantified by scintillation counting.

- 5 In a typical experiment basal activity is 300 dpm per 75 μ l sample of reagent controls, which is increased to 3000 dpm in the presence of enzyme. Compound activity is expressed as IC_{50} (the concentration of drug substance which gives 50% enzyme inhibition in the assay) and L-NMMA, which gives an IC_{50} of about 0.4 μ M is tested as a standard to verify the procedure. Compounds are tested at a range of concentrations and from the
- 10 inhibitions obtained IC_{50} values are calculated.

In this screen the compounds of Examples 1 to 30 give IC_{50} values less than 25 μ M, indicating that they are predicted to show useful therapeutic activity.

CLAIMS:

1. A compound of formula (I)



wherein

R^1 and R^2 independently represent hydrogen, C1 to 6 alkyl, C2 to 6 alkenyl, C2 to 6
 10 alkynyl, C1 to 6 alkoxy, C1 to 6 alkylthio, halogen, hydroxy, trifluoromethyl or amino;

R^3 represents one or more substituents independently selected from hydrogen, C1 to 6
 alkyl, C2 to 6 alkenyl, C2 to 6 alkynyl, C1 to 6 alkoxy, C1 to 6 alkylthio, halogen,
 hydroxy, trifluoromethyl, amino, cyano, nitro, trifluoromethoxy, methanesulphonyl,
 15 sulphonyl, $-NR^4R^5$, $-COOR^6$, $-CONR^7R^8$, benzyloxy, phenyl, or a 5-membered
 heterocyclic aromatic ring containing 1 to 3 heteroatoms which may be the same or
 different and are selected from O, N and S, which phenyl or 5-membered heterocyclic
 aromatic ring is optionally substituted, the optional substituents being C1 to 6 alkyl,
 halogen, cyano, nitro, hydroxy, C1 to 6 alkoxy, trifluoromethyl and trifluoromethoxy;

20

R^4 , R^5 and R^6 independently represent hydrogen or C1 to 6 alkyl;

R^7 and R^8 independently represent hydrogen, C1 to 6 alkyl or phenyl, which phenyl is
 optionally substituted by one or more groups independently selected from C1 to 6 alkyl,
 25 halogen, cyano, nitro, hydroxy, C1 to 6 alkoxy, trifluoromethyl and trifluoromethoxy;

X represents $-(CH_2)_n-$, wherein n represents zero or 1; and

A represents an aromatic carbocyclic ring or a 5- or 6-membered heterocyclic aromatic ring
5 containing 1 to 3 heteroatoms which may be the same or different and are selected from O,
N and S;

or a pharmaceutically acceptable salt, enantiomer, racemate or tautomer thereof.

10 2. A compound of formula (I), according to Claim 1, wherein A represents a benzo ring.

3. A compound of formula (I), according to Claim 1 or Claim 2, wherein R^1 represents
hydrogen.

15 4. A compound of formula (I), according to Claim 1 or Claim 2, wherein R^1 represents
fluoro.

5. A compound of formula (I), according to any one of Claims 1 to 4, wherein R^2
represents fluoro.

20

6. A compound of formula (I), according to any one of Claims 1 to 5, wherein X
represents $-(CH_2)_n-$ and n represents zero.

7. A compound of formula (I) which is:

25

(R^*R^*)-4'-amino-5'-fluoro-1,3,4,10a-tetrahydrospiro[pyrido[2,1-a]isoindole-
2,2'(1'H)quinazoline]-6-one;
(R^*S^*)-4'-amino-5'-fluoro-1,3,4,10a-tetrahydrospiro[pyrido[2,1-a]isoindole-
2,2'(1'H)quinazoline]-6-one;

- (R R*)-4-amino-5-fluoro-3b',4',6',7'-tetrahydrospiro[quinazoline-2,5'(2H,9'H)-thieno[3,2-a]indolizine]-9'-one;
- (R*S*)-4-amino-5-fluoro-3b',4',6',7'-tetrahydrospiro[quinazoline-2,5'(2H,9'H)-thieno[3,2-a]indolizine]-9'-one;
- 5 (R*R*)-4-amino-5-fluoro-6',7',9',9a'-tetrahydrospiro[quinazoline-2,8'(2H,4'H)-thieno[2,3-a]indolizine]-4'-one;
- (R*S*)-4-amino-5-fluoro-6',7',9',9a'-tetrahydrospiro[quinazoline-2,8'(2H,4'H)-thieno[2,3-a]indolizine]-4'-one;
- (R*R*)-4'-amino-7-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-10 2,2'(1'H)-quinazoline]-6-one;
- (R*S*)-4'-amino-7-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (R*R*)-4'-amino-8-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- 15 (R*S*)-4'-amino-8-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (R*R*)-4'-amino-9-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (R*S*)-4'-amino-9-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-20 2,2'(1'H)-quinazoline]-6-one;
- (R*R*)-4'-amino-10-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (R*S*)-4'-amino-10-chloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- 25 (R*R*)-4'-amino-7,8-dichloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (R*S*)-4'-amino-7,8-dichloro-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (R*R*)-4'-amino-7-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-30 2,2'(1'H)-quinazoline]-6-one;

- (*R S*)-4'-amino-7-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R*R**)-4'-amino-8-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- 5 (*R*S**)-4'-amino-8-methyl-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R*R**)-4'-amino-8-methoxy-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R*S**)-4'-amino-8-methoxy-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-10 2,2'(1'H)-quinazoline]-6-one;
- (*R*R**)-4'-amino-8-cyano-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R*S**)-4'-amino-8-cyano-5'-fluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- 15 (*R*R**)-4'-amino-8-cyano-5',8'-difluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R*S**)-4'-amino-8-cyano-5',8'-difluoro-1,3,4,10b-tetrahydrospiro[pyrido[2,1-a]isoindole-2,2'(1'H)-quinazoline]-6-one;
- (*R*R**)-4'-amino-5'-fluoro-7,8,10,10a-tetrahydrospiro[pyrido[3,4-a]indolizine-9,2'(1'H)-20 quinazoline]-5-one;
- (*R*S**)-4'-amino-5'-fluoro-7,8,10,10a-tetrahydrospiro[pyrido[3,4-a]indolizine-9,2'(1'H)-quinazoline]-5-one;
- (*R*R**)-4'-amino-5'-fluoro-6b,7,9,10-tetrahydrospiro[naphtho[2,1-a]indolizine-8,2'(1'H)-quinazoline]-12-one;
- 25 (*R*S**)-4'-amino-5'-fluoro-6b,7,9,10-tetrahydrospiro[naphtho[2,1-a]indolizine-8,2'(1'H)-quinazoline]-12-one;

or a pharmaceutically acceptable salt, enantiomer or tautomer thereof.

8. A compound of formula (I), according to any one of Claims 1 to 7, or a
- 30 pharmaceutically acceptable salt, enantiomer or tautomer thereof, for use as a medicament.

9. A pharmaceutical composition comprising a compound of formula (I) according to any one of Claims 1 to 7, or a pharmaceutically acceptable salt, enantiomer or tautomer thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier.

10. The use of a compound of formula (I) according to any one of Claims 1 to 7, or a pharmaceutically acceptable salt, enantiomer or tautomer thereof, in the manufacture of a medicament for the treatment or prophylaxis of human diseases or conditions in which inhibition of nitric oxide synthase activity is beneficial.

11. The use as claimed in Claim 10 wherein it is predominantly inducible nitric oxide synthase that is inhibited.

12. The use of a compound of formula (I) as defined in any one of Claims 1 to 7, or a pharmaceutically acceptable salt, enantiomer or tautomer thereof, in the manufacture of a medicament, for the treatment or prophylaxis of inflammatory diseases.

13. The use as claimed in Claim 12 wherein the disease is asthma or rheumatoid arthritis.

14. The use of a compound of formula (I) as defined in any one of Claims 1 to 7, or a pharmaceutically acceptable salt, enantiomer or tautomer thereof, in the manufacture of a medicament, for the treatment or prophylaxis of pain.

15. The use of a compound of formula (I) as defined in any one of Claims 1 to 7, or a pharmaceutically acceptable salt, enantiomer or tautomer thereof, in combination with a COX-2 inhibitor, in the manufacture of a medicament, for the treatment or prophylaxis of inflammatory diseases.

16. A method of treating, or reducing the risk of, human diseases or conditions in which inhibition of nitric oxide synthase activity is beneficial which comprises administering a therapeutically effective amount of a compound of formula (I), as defined in any one of

Claims 1 to 7, or a pharmaceutically acceptable salt, enantiomer or tautomer thereof, to a person suffering from, or at increased risk of, such diseases or conditions.

17. A method of treatment according to Claim 16 in which it is predominantly inducible
5 nitric oxide synthase that is inhibited.

18. A method of treating, or reducing the risk of, inflammatory disease in a person suffering from, or at risk of, said disease, wherein the method comprises administering to the person a therapeutically effective amount of a compound of formula (I), as defined in any one of
10 Claims 1 to 7, or a pharmaceutically acceptable salt, enantiomer or tautomer thereof.


19. The method of treatment as claimed in Claim 18 wherein the disease is asthma or rheumatoid arthritis.

15 20. A method of treating, or reducing the risk of, pain in a person suffering from, or at risk of, said condition, wherein the method comprises administering to the person a therapeutically effective amount of a compound of formula (I), as defined in any one of Claims 1 to 7, or a pharmaceutically acceptable salt, enantiomer or tautomer thereof.

20 21. A method of treating, or reducing the risk of, inflammatory disease in a person suffering from, or at risk of, said disease, wherein the method comprises administering to the person a therapeutically effective amount of a combination of a compound of formula (I), as defined in any one of Claims 1 to 7, or a pharmaceutically acceptable salt, enantiomer or tautomer thereof, with a COX-2 inhibitor.


25

22. A compound of formula (III)




wherein R^3 , A and X are as defined in Claim 1,
with the proviso that R^3 does not represent hydrogen.

23. A process for the preparation of a compound of formula (I), as defined in any one of
5 Claims 1 to 7, or a pharmaceutically acceptable salt, enantiomer or tautomer thereof,
wherein the process comprises reaction of a compound of formula (II), or a salt thereof:



- wherein R^1 and R^2 are as defined in Claim 1,
10 with a compound of formula (III) or a protected derivative thereof:



wherein R^3 , A and X are as defined in claim 1.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 98/01206

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 487/20, C07D 471/20, C07D 471/04, A61K 31/505, A61K 31/445
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D

Documentation searched other than minimum: documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9714686 A1 (ASTRA PHARMACEUTICALS LIMITED), 24 April 1997 (24.04.97)	1-23
	--	
A	Tetrahedron, Volume 49, No 11, 1993, Jean-Pierre Gesson et al, "Alpha-N-Acyliminium Ion - 2-Bromoalkene Cyclizations", page 2239 - page 2248, page 2243, compound 11	22
	--	
A	STN International, File CAPLUS, CAPLUS accession no. 1985:95488, Lochead, Alistair W. et al: "Use of chloroalkenylamines for the synthesis of 1-azabicyclo(3.3.0)octane and 1-azabicyclo (4.3.0)nonane derivatives", J. Chem. Soc., Perkin Trans. 1 (1984), (11), 2477-89	1-22
	-- -----	

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to underpin the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

30 Sept 1998

Date of mailing of the international search report

16-10-1998

Name and mailing address of the ISA

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Gerd Strandell

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 98/01206

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16-21
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 16-21 relate to methods of treatment of the human or animal body by surgery or by therapy. See PCT, Rule 39.1(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/SE 98/01206

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9714686 A1	24/04/97	AU 7224396 A	07/05/97
		EP 0858451 A	19/08/98
		GB 9521231 D	00/00/00
		AU 7321796 A	22/05/97
		EP 0854996 A	29/07/98
		GB 9602668 D	00/00/00
		GB 9614386 D	00/00/00
<hr/>			